

MAGNETIC PROPELLER OUTFLOWS

R.V.E. LOVELACE

Department of Astronomy, Cornell University, Ithaca, NY 14853-6801; rvl1@cornell.edu

M.M. ROMANOVA

Space Research Institute, Russian Academy of Sciences, Moscow, Russia; and
Department of Astronomy, Cornell University, Ithaca, NY 14853-6801; romanova@astro.sun.tn.cornell.edu

G.S. BISNOVATYI-KOGAN

Space Research Institute, Russian Academy of Sciences, Moscow, Russia; gkogan@mx.iki.rssi.ru

Accepted to the Astrophysical Journal

ABSTRACT

A model is developed for magnetic ‘propeller’-driven outflows which cause a rapidly rotating magnetized star accreting from a disk to spin-down. Energy and angular momentum lost by the star goes into expelling most of the accreting disk matter. The theory gives an expression for the effective Alfvén radius R_A (where the inflowing matter is effectively stopped) which depends on the mass accretion rate, the star’s mass and magnetic moment, *and* the star’s rotation rate. The model points to a mechanism for ‘jumps’ between spin-down and spin-up evolution and for the reverse transition, which are changes between two possible equilibrium configurations of the system. In for example the transition from spin-down to spin-up states the Alfvén radius R_A decreases from a value larger than the corotation radius to one which is smaller. In this transition the ‘propeller’ goes from being “on” to being “off.” The ratio of the spin-down to spin-up torque (or the ratio for the reverse change) in a jump is shown to be of order unity.

Subject headings: accretion, accretion disks—plasmas—magnetic fields—stars: magnetic fields—X-rays: stars

1. INTRODUCTION

Observations of some X-ray pulsars show remarkable ‘jumps’ between states where the pulsar is spin-ning-down to one where it is spinning-up. Examples include the objects Cen X-3 (Chakrabarty et al. 1993) and GX 1+4 (Chakrabarty et al. 1997; Cui 1997). The theoretical problem of disk accretion to a rotating magnetized star has been discussed in many works over a long period (Pringle & Rees 1972; Lynden-Bell & Pringle 1974; Ghosh & Lamb 1979; Wang 1979; Lipunov 1993; Shu et al. 1994; Lovelace, Romanova, & Bisnovaty-Kogan 1995 (hereafter LRBK); Li & Wickramasinghe 1997). However, except for the work by Li & Wickramasinghe (1997), the studies do not specifically address the ‘propeller’ regime (Illarionov & Sunyaev 1975) where the rapid rotation of the star’s magnetosphere acts to expell most of the accreting matter and where the star spins-down. Recent computer simulation studies of disk accretion to a rotating star with an aligned dipole magnetic field (Hayashi, Shibata, & Matsumoto 1996; Goodson, Winglee, & Böhm 1997; Miller & Stone 1997) provide evidence of time-dependent outflows but do not give definite evidence for a ‘propeller’ regime with spin-down of the star. The present work considers the ‘propeller’ regime and develops a simple physical model where the energy and angular momentum lost by the rotating star goes into a magnetically driven outflow.

2. THEORY

We consider the problem of disk accretion onto a rotating magnetized star which has an aligned dipole magnetic

field. We focus on the limit where the star is rotating rapidly and the disk-star configuration is as sketched in Figure 1 (see Figure 3 of LRBK). We consider the flow of mass, angular momentum, and energy into and out of the annular region indicated by the box $A'ACC'$ in this figure, where $A'A$ is at radius r_1 and CC' is at r_2 . Notice that at this point the values of r_1 and r_2 are *unknown*. They are determined by the physical considerations discussed here.

Consider first the outer surface CC' through the disk. The influx of mass into the considered region is

$$\dot{M}_{accr} = 2\pi r_2 \int_{-h}^h dz [\rho (-v_r)]_2 > 0, \quad (1)$$

where h is the half-thickness of the disk, and the 2 subscript indicates evaluation at $r = r_2$. We assume that mass accretion rate for $r \geq r_2$ is approximately constant equal to $\dot{M}_{accr}$. That is, we consider that outflow from the disk is negligible for $r \geq r_2$. The influx of angular momentum into the considered region is

$$\dot{L}_{accr} = \dot{M}_{accr} (rv_\phi)_2 - T_2. \quad (2a)$$

Here,

$$T_2 = 2\pi r_2^2 \int_{-h}^h dz (T_{r\phi}^{vis})_2, \quad (2b)$$

where $T_{r\phi}^{vis}$ is the viscous contribution to the stress tensor which includes *both* the turbulent hydrodynamic and turbulent magnetic stresses. The influx of energy into the considered region is

$$\dot{E}_{accr} = \dot{M}_{accr} \left(-\frac{GM}{2r_2} + w_2 \right) - \omega_2 T_2, \quad (3)$$




FIG. 1.— Geometry of disk accretion to a rapidly rotating star with an aligned dipole magnetic field. Here, r_1 and r_2 are the boundaries of the region considered in the text; R_A is the effective radius where the outflow begins; ω_* is the star's rotation rate; ω_K is the Keplerian rotation rate of the accretion disk; and $r_{cr} \equiv (GM/\omega_*^2)^{1/3}$ is the corotation radius. The magnetic field in the vicinity of r_1 has an essential time-dependence owing to the continual processes of stellar flux leaking outward into the disk, the resulting field loops being inflated by the differential rotation (LRBK), and the reconnection between the open disk field and the closed stellar field loops. The dashed lines marked by the letters 'n' indicate neutral surfaces along which reconnection occurs.

where $\omega \equiv v_\phi/r \approx v_K/r$ with $v_K \equiv \sqrt{GM/r}$ the Keplerian speed and M the mass of the star, and where w_2 is the enthalpy. For conditions of interest here the disk at r_2 is geometrically thin so that $w \sim c_s^2 \ll v_K^2$, where c_s is the sound speed.

Consider next the fluxes of mass, angular momentum, and energy across the surface AA' in Figure 1. For the physical regime considered, where $v_K(r_1)/r_1 \ll \omega_*$ ($=$ the angular rotation rate of the star), the mass accretion across the AA' surface is assumed to be small compared with $\dot{M}_{accr}$. The reason for this is that any plasma which crosses the AA' surface will be 'spun-up' to an angular velocity ω_* (by the magnetic force) which is substantially larger than the Keplerian value, and thus it will be thrown outwards. Thus the efflux of angular momentum across this surface from the considered region is $\dot{L}_1 = -T_1$. The efflux of energy across this surface is $\dot{E}_1 = -\omega_* T_1$, where ω_* is the angular rotation rate of the star and the inner magnetosphere as shown in Figure 1. For the conditions considered, the star slows down and loses rotational energy so that $T_1 > 0$.

We have $\dot{E}_1/\dot{L}_1 = \omega_*$. Because the interaction of the star with the accretion flow is by assumption entirely across the surface AA' , this is consistent with the spin-down of a star with constant moment of inertia I ; that is, $\Delta E_*/\Delta L_* = I_* \omega_* \Delta \omega_*/(I_* \Delta \omega_*) = \omega_*$.

Next we consider the mass, angular momentum, and energy fluxes across the surfaces $A'C'$ and AC in Figure 1.

As mentioned, accretion to the star is small for $r_{cr} \ll r_1$ where r_{cr} is the corotation radius as indicated in Figure 1. Thus, the mass accretion goes mainly into outflows, $\dot{M}_{out} \approx \dot{M}_{accr}$, where "out" stands for outflows. The angular momentum outflow across the surfaces $A'C'$ and AC , $\dot{L}_{out}$, must be the difference between the angular momentum lost by the star and the incoming angular momentum of the accretion flow. The angular momentum carried by radiation from the disk is negligible because $(v_K/c)^2 \ll 1$, where c is the speed of light. That is, $\dot{L}_{out} = \dot{L}_{accr} - \dot{L}_1$. The energy outflow across the $A'C'$ and AC surfaces is $\dot{E}_{out} + \dot{E}_{rad} = \dot{E}_{accr} - \dot{E}_1$, where $\dot{E}_{rad}$ is the radiation energy loss rate from the disk surfaces between $r = r_1$ and r_2 , and $\dot{E}_{out}$ is the rate of energy loss carried by the outflows.

Angular momentum conservation gives

$$\dot{L}_{out} = \dot{M}_{accr}(rv_\phi)_2 - T_2 + T_1. \quad (4)$$

We have

$$\dot{E}_{out} + \dot{E}_{rad} = -\frac{GM\dot{M}_{accr}}{2r_2} - \omega_2 T_2 + \omega_* T_1. \quad (5)$$

We can solve equation (4) for T_2 and thereby eliminate this quantity from the energy equation (5). This gives

$$\dot{E}_{out} + \dot{E}_{rad} = -\frac{3GM\dot{M}_{accr}}{2r_2} + \omega_2 \dot{L}_{out} + (\omega_* - \omega_2) T_1. \quad (6)$$

The preceeding equations are independent of the nature of the outflows from the disk. At this point we consider the case of magnetically driven outflows as treated by Lovelace, Berk, and Contopoulos (1991, hereafter LBC). In the LBC model the outflows come predominantly from an annular inner region of the disk of radius $\sim R_A$ where the disk rotation rate is $\omega_0 = \sqrt{GM/R_A^3}$. Thus we assume that the outflows come from a region of the disk which is approximately in Keplerian rotation. For the present situation, shown in Figure 1, it is clear that we must have $r_1 < R_A < r_2$. Further, we will assume R_A is close in value to r_1 with $(R_A - r_1)/R_A \lesssim 1$. For conditions where the outflow from the disk is relatively low temperature (sound speed much less than Keplerian speed), equations (16) and (18) of LBC imply the general relation $\dot{E}_{out} = \omega_0 \dot{L}_{out} - 3GM\dot{M}_{out}/(2R_A)$. This equation can be used to eliminate $\dot{L}_{out}$ in favor of $\dot{E}_{out}$ in equation (6). Recalling that $\dot{M}_{out} = \dot{M}_{accr}$ we have

$$(1 - \delta^{\frac{3}{2}})\dot{E}_{out} + \dot{E}_{rad} = (\omega_* - \omega_2)T_1 - \frac{3GM\dot{M}_{accr}}{2r_2}(1 - \delta^{\frac{1}{2}}), \quad (7)$$

where $\delta \equiv R_A/r_2 < 1$.

The energy dissipation in the region of the disk $r = r_1$ to r_2 heats the disk and this heat energy is transformed into outgoing radiation $\dot{E}_{rad}$. Thus we have

$$\dot{E}_{rad} = \frac{3}{2} \int_{r_1}^{r_2} dr \frac{GM\dot{M}_{disk}(r)}{r^2}, \quad (8)$$

(Shakura 1973; Shakura & Sunyaev 1973), where $\dot{M}_{disk}(r_2) = \dot{M}_{accr}$ and $\dot{M}_{disk}(r_1) \approx 0$. The essential change in $\dot{M}_{disk}(r)$ occurs in the vicinity of R_A so that $\dot{E}_{rad} \approx (3GM\dot{M}_{accr}/2)(R_A^{-1} - r_2^{-1})$. Thus equation (7) becomes

$$(1 - \delta^{\frac{3}{2}})\dot{E}_{out} = (\omega_* - \omega_2)T_1 - \frac{3GM\dot{M}_{accr}}{2R_A}(1 - \delta^{\frac{3}{2}}). \quad (9)$$

Thus the power from the spin-down of the star $\omega_* T_1$ must be larger than a certain value in order to drive the outflow.

For magnetically driven outflows, the value of $\dot{E}_{out}$ can be written as

$$\dot{E}_{out} = \frac{3}{2} \mathcal{F}_0^{\frac{2}{3}} \omega_0^{\frac{4}{3}} R_A^{\frac{8}{3}} B_0^{\frac{4}{3}} \dot{M}_{accr}^{\frac{1}{3}}, \quad (10)$$

(equation 34 of LBC), where $\mathcal{F}_0$ is a dimensionless numerical constant $\gtrsim 0.234$, and B_0 is the poloidal magnetic field at the base of the outflow at $r = R_A$. We take the simple estimate $B_0 = \mu/R_A^3$, which omits corrections for example for compression of the star's field by the inflowing plasma.

Next we consider the torque on the star T_1 . Because most of the matter inflowing in the accretion disk at $r = r_2$ is driven off in outflows at distances $r > r_1$, the stress is necessarily due to the magnetic field. The magnetic field in the vicinity of r_1 has an essential time-dependence owing to the continual processes of stellar flux leaking outward into the disk, the resulting field loops being inflated by the differential rotation (LRBK), and the

reconnection between the open disk field and the closed stellar field loops. The time scale of these processes is $t_1 \lesssim 2\pi r_1/v_K(r_1)$. We make the estimate of the torque $T_1 = -2\pi r_1^2(2\Delta z)\langle B_r B_\phi \rangle_1/(4\pi)$, where Δz is the vertical half-thickness of the region where the magnetic stress is significant, and where the angular brackets denote a time average of the field quantities at $r \sim r_1$. The magnetic field components, B_r , B_ϕ , with $B_\phi \propto -B_r$ necessarily, must be of magnitude less than or of the order of the dipole field $B_1 = \mu/r_1^3$ at $r = r_1$. The fact that B_ϕ has the opposite sign to that of B_r is due to the fact that the B_ϕ field arises differential rotation between the region $r < r_1$, which rotates at rate ω_* , and the region $r > r_1$, which rotates at rate $\omega_K(r_1) < \omega_*$. Also, it is reasonable to assume $\Delta z \lesssim r_1$. Therefore,

$$T_1 \approx \bar{\alpha} r_1^3 B_1^2 \equiv \bar{\alpha} \frac{\mu^2}{R_A^3}, \quad (11)$$

where $\bar{\alpha} \leq 1$ (the time average of $\alpha(t)$) is a dimensionless constant analogous to the α -parameter of Shakura (1973) and Shakura & Sunyaev (1973). (Note that because $r_1 \lesssim R_A$, $\bar{\alpha}$ and $\bar{\alpha}_1 = (r_1/R_A)^3 \bar{\alpha}$ are of the same order of magnitude.)

Substituting equations (10) and (11) into equation (9) gives

$$\frac{3}{2} (1 - \delta^{\frac{3}{2}}) \mathcal{F}_0^{\frac{2}{3}} \left(\frac{r_A}{R_A} \right)^{\frac{7}{3}} = \frac{\bar{\alpha} r_A^{\frac{7}{3}}}{R_A^2 r_{cr}^{\frac{3}{3}}} \left[1 - \left(\frac{r_{cr}}{R_A} \right)^{\frac{3}{2}} \delta^{\frac{3}{2}} \right] - \frac{3}{2} (1 - \delta^{\frac{3}{2}}). \quad (12)$$

Here, we have introduced two characteristic radii – the corotation radius,

$$r_{cr} \equiv \left(\frac{GM}{\omega_*^2} \right)^{1/3} \approx 1.5 \times 10^8 \text{ cm } M_1^{1/3} P_1^{2/3},$$

with $P_1 \equiv (2\pi/\omega_*)/1\text{s}$ the pulsar period and $M_1 \equiv M/M_\odot$. [For a young stellar object, $r_{cr} \approx 1.36 \times 10^{12} M_1^{1/3} P_{10d}^{2/3}$, where P_{10d} is the period in units of 10 days.] The second is the *nominal* Alfvén radius

$$r_A \equiv \left[\frac{\mu^2}{\dot{M}_{accr} \sqrt{GM}} \right]^{2/7} \approx 3.6 \times 10^8 \text{ cm } \frac{\mu_{30}^{4/7}}{\dot{M}_{17}^{2/7} M_1^{1/7}},$$

where the accretion rate $\dot{M}_{17} \equiv \dot{M}_{accr}/(10^{17} \text{ g/s})$ with $10^{17} \text{ g/s} \approx 1.6 \times 10^{-9} M_\odot/\text{yr}$ and $\mu_{30} \equiv \mu/(10^{30} \text{ Gcm}^2)$ with the magnetic field at the star's equatorial surface $\mu/r^3 = 10^{12} G(10^6 \text{ cm}/r)^3$. [For a young stellar object $r_A \approx 1.81 \times 10^{12} \text{ cm } \mu_{36.5}^{4/7} / (\dot{M}^{2/7} M_1^{1/7})$, where the normalization corresponds to a stellar radius of 10^{11} cm , a surface magnetic field of $3 \times 10^3 \text{ G}$, and an accretion rate of $1.6 \times 10^{-8} M_\odot/\text{yr}$.] The corotation radius is the distance from the star where the centrifugal force on a particle corotating with the star ($\omega_*^2 r$) balances the gravitational attraction (GM/r^2). The Alfvén radius r_A is the distance from a *non-rotating* star where the free-fall of a quasi-spherical accretion flow is stopped, which occurs (approximately) where the kinetic energy-density of the flow equals




FIG. 2.— The plot shows the *effective* Alfvén radius R_A (normalized by the *nominal* Alfvén radius r_A) as a function of the normalized corotation radius r_{cr}/r_A for $\bar{\alpha} = 0.2$ obtained from equation (12). We have neglected the $\delta^{3/2} = (R_A/r_2)^{3/2}$ terms compared with unity and taken the value $\mathcal{F}_0 = 0.234$ from LBC. Only the part of the solid curve above the dashed line $R_A = r_{cr}$ is consistent with our assumptions. The dashed horizontal line r_{to} indicates the “turnover radius” of the disk rotation curve or effective Alfvén radius in the regime discussed by LRBK where $\dot{M}_{accr}$ falls onto the star and the star spins-up. For this line the turbulent magnetic diffusivity of the disk is taken to be $\alpha D = 0.1$. The points a b c d and the associated vertical lines represent possible transitions between spin-down and spin-up of the pulsar as discussed in the text.

the energy-density of the star’s dipole field. Note that the assumptions leading to equation (12) require $R_A > r_{cr}$.

Notice that R_A (or $r_1 \sim R_A$) is the *effective* Alfvén radius for a *rotating* star. It depends on *both* r_A and r_{cr} in contrast with the common notion that the Alfvén radius is given by r_A even for a rotating star. From equation (11), the *spin-down* rate of the star is $I(d\omega_*/dt) = -\bar{\alpha} \mu^2/R_A^3$, where I is the moment of inertia of the star (assumed constant). Thus the spin-down rate depends on both r_A and r_{cr} .

Figure 2 shows the dependence of R_A on r_A and r_{cr} for a sample case. For conditions of a newly formed disk around a young pulsar, the initial system point would be on the upper left-hand part of the curve. Due to the pulsar slowing down (assuming μ and $\dot{M}$ constant), the system point would move downward and to the right as indicated by the arrow. In this region of the diagram, $R_A \approx \sqrt{\bar{\alpha}} r_A^{7/4}/r_{cr}^{3/4} \propto \sqrt{\omega}$ (for $\delta \ll 1$), so that the torque on the star is $T_1 \approx (GMM)^{3/2}/(\mu\sqrt{\bar{\alpha}} \omega^{3/2})$. Thus the braking index is $n = -3/2$, where n is defined by the relation $\dot{\omega}_* = -\text{const } \omega_*^n$. Numerically,

$$\frac{1}{P} \frac{dP}{dt} = \frac{1.55 \times 10^{-5}}{\text{yr}} \frac{P_1^{5/2} (M_1 \dot{M}_{17})^{3/2}}{\sqrt{\bar{\alpha}} \mu_{30} I_{45}}, \quad (13)$$

where $I_{45} \equiv I/(10^{45} \text{ g cm}^2)$. For $R_A \gg r_{cr}$, the mass accretion rate to the star is small compared with $\dot{M}_{accr}$, but some accretion may occur due to ‘leakage’ of relatively low angular momentum plasma across field lines near r_1 (Arons & Lea 1976).

As R_A decreases, the spin-down torque on the star increases. Over a long interval, R_A will decrease to a value larger than r_{cr} but not much larger. In this limit, mass accretion to the star $\dot{M}_1$ may become significant. Our treatment can be extended to this limit by noting that $\dot{M}_1 = \dot{M}_2 - \dot{M}_{out}$, $\dot{L}_1 = \dot{M}_1 \omega_1 r_1^2 - T_1$, and $\dot{E}_1 = \dot{M}_1 (-GM/2r_1) - \omega_* T_1$, where T_1 is given by equation (11). In this limit the accretion luminosity is GMM_1/r_* , where r_* is the star’s radius. Figure 2 is not changed appreciably for $\dot{M}_1 < \dot{M}_{accr}$.

Further spin-down of the star will cause the system point in Figure 2 to approach the right-most part of the curve. Further spin-down of the star is impossible. At this point of the evolution, the only possibility is a transition to the spin-up regime. In this regime the effective Alfvén radius is the ‘turnover radius’ r_{to} of the disk rotation curve calculated by LRBK, the star spins-up at the rate $I d\omega_*/dt = \dot{M}_{accr} (GM r_{to})^{1/2}$, and most of the disk accretion $\dot{M}_{accr}$ falls onto the star. The dashed horizontal line in Figure 2 indicates r_{to} which is necessarily less than r_{cr} .

The location of the turnover line r_{to} in Figure 2 suggests the possible evolution shown by the sequence of points $a \rightarrow b \rightarrow c \rightarrow d \rightarrow a$. The system can jump down from point a where the star spins-down to point b where it spins-up. The spin-down torque at a is $-\bar{\alpha} \mu^2/R_{Aa}^3$ (where R_{Aa} is the effective Alfvén radius at point a), whereas the spin-up torque at b is $\dot{M}_{accr} (GM r_{to})^{1/2}$. The magnitude of the ratio of these torques is

$$\frac{\text{spin-down}}{\text{spin-up}} = \bar{\alpha} \left(\frac{r_A}{R_{Aa}} \right)^{7/2} \left(\frac{R_{Aa}}{r_{to}} \right)^{1/2}. \quad (14)$$

With the system on the r_{to} line, it evolves to the left. Because $r_{to} \leq r_{cr}$, there must be an upward jump from point c to point d . For this case the torque ratio is given by equation (14) with $R_{Aa} \rightarrow R_{Ad}$. From point d the system evolves to the right. For the example shown in Figure 2, the torque ratio is ≈ 3.45 for $a \rightarrow b$ whereas it is ≈ 0.87 for $c \rightarrow d$. The vertical line $c \rightarrow d$ is at the left-most position allowed for the considered conditions, but this transition could also occur if the line is shifted to the right. The line $a \rightarrow b$ can be displaced slightly to the right or it can be displaced to the left to be coincident with the $c \rightarrow d$ line. In the latter case the torque ratio for the spin-down to spin-up jump is approximately equal to the torque ratio for the spin-up to spin-down jump and is ≈ 0.87 . The smaller the horizontal separation of the $a \rightarrow b$ and the $c \rightarrow d$ lines, the shorter is the time interval between jumps.

Summarizing, we can say that the horizontal locations of the transitions, $a \rightarrow b$, and $c \rightarrow d$, are indeterminate

within a definite range. The locations of the jumps in the (R_A, r_{cr}) plane may in fact be a stochastic or chaotic in nature and give rise to chaotic hysteresis in the of the spin-down/spin-up behavior of the pulsar. The jumps could be triggered by small variations in the accretion flow ($\dot{M}$ for example) and magnetic field configuration (the time-dependence of α in the torque T_1). Analysis of the accreting neutron star system Her X-1 (Voges, Atmanspacher, & Scheingraber 1987; Morfill et al. 1989) suggests that the intensity variations are described by a low dimensional deterministic chaotic model. The transistions between spin-down and spin-up and the reverse transistions may be described by an analogous model.

The allowed values in Figure 2 have $r_{cr}/r_A \leq \text{const} \equiv k$, where $k \approx \bar{\alpha}^{2/3}$. This corresponds to pulsar periods

$$P \leq 5.2\text{s} \left(\frac{\bar{\alpha}}{M_1^{5/7}} \right) \left(\frac{\mu_{30}^2}{\dot{M}_{17}} \right)^{3/7}. \quad (15)$$

For some long period pulsars such as GX 1+4 this inequality points to magnetic moment values μ_{30} appreciably larger than unity. Periods much longer than allowed by (15) can result for pulsars which accrete from a stellar wind (Bisnovaty-Kogan 1991). [For a young stellar object, equation (15) gives $P \leq 8\text{d}(\bar{\alpha}/M_1^{5/7})(\mu_{36.5}^2/\dot{M}_{18})^{3/7}$.]

3. DISCUSSION

This work presents a new investigation of the ‘propeller’ regime of disk accretion to a rapidly rotating magnetized star. The work considers the field configuration proposed by LRBK, the theory of LBC on magnetically driven jets, and the conservation of mass, angular momentum, and

energy to derive an expression for the effective Alfvén radius R_A (equation 12) and the spin-down torque on the star T_1 (equation 11). Our work is in qualitative accord with that of Li and Wickramasinghe (1997) who also consider the propeller effect of Illarionov and Sunyaev (1975). Our Figure 1 is similar to Figure 4 of Li and Wickramasinghe for the spin-down regime, and our earlier work on the spin-up regime (LRBK) agrees with their Figure 3. We find that R_A depends not only on μ , $\dot{M}$, and M , but also on the star’s rotation rate ω_* . Because R_A decreases as ω_* decreases, there is a minimum value of ω_* or a maximum value of the pulsar period $P = 2\pi/\omega_*$. The model points to a mechanism for ‘jumps’ between spin-down and spin-up evolution (and the reverse transition). In our picture, in a spin-down to spin-up transition, for example, the effective Alfvén radius decreases by an appreciable factor going from $> r_{cr}$ to $< r_{cr}$. The propeller goes from being “on” to being “off” in this transistion, which is a change between two possible equilibrium configurations. The transistions may be stochastic or chaotic in nature with triggering due to small variations in the accretion flow or in the magnetic field configuration. The ratio of the spin-down to spin-up torques (or the ratio for the reverse transition) is found to be of order unity (equation 14). This agrees with observations of for example Cen X-3 (Chakrabarty et al. 1993) and GX 1+4 (Chakrabarty et al. 1997; Cui 1997).

We thank Dr. Wei Cui for valuable comments on this work. This work was supported in part by NSF grant AST-9320068. Also, this work was made possible in part by Grant No. RP1-173 of the U.S. Civilian R&D Foundation for the Independent States of the Former Soviet Union. The work of RVEL was also supported in part by NASA grant NAG5 6311.

REFERENCES

- Arons, J., & Lea, S.M. 1976, ApJ, 207, 914
 Bisnovaty-Kogan, G.S. 1991, A&A, 245, 528
 Chakrabarty, D. et al. 1993, ApJ, 403, L33
 Chakrabarty, D. et al. 1997, ApJ, 481, L105
 Cui, W. 1997, ApJ, 482, L163
 Ghosh, P., & Lamb, F.K. 1979, ApJ, 234, 296
 Goodson, A.P., Winglee, R.M., & Böhm, K.H. 1997, ApJ, 489, 390
 Hayashi, M.R., Shibata, K., & Matsumoto, R. 1996, ApJ, 468, L37
 Illarionov, A.F., & Sunyaev, R.A. 1975, A&A, 39, 185
 Li, J., & Wickramasinghe, D.T. 1997, in *Accretion Phenomena and Related Outflows, IAU Colloquium 163*, ASP Conf. Ser., Vol. 121, 241
 Lipunov, V.M. 1993, *Astrophysics of Neutron Stars*, (Springer-Verlag: Berlin)
 Lovelace, R.V.E., Berk, H.L., & Contopoulos, J. 1991, ApJ, 379, 696
 Lovelace, R.V.E., Romanova, M.M., & Bisnovaty-Kogan, G.S. 1995, MNRAS, 275, 244
 Lynden-Bell, D., & Pringle, J.E. 1974, MNRAS, 168, 603
 Miller, K., & Stone, J. 1997, ApJ, 489, 890
 Morfill, G.E., Atmanspacher, H., Demmel, V., Scheingraber, H., & Voges, W. 1989, in *Timing Neutrons Stars*, H. Ögelman & E.P.J. van den Heuvel, eds. (Kluwer Acad. Pub.), p. 71
 Pringle, J.E., & Rees, M.J. 1972, A&A, 21, 1
 Shakura, N.I. 1973, SvA, 16, 756
 Shakura, N.I., & Sunyaev, R.A. 1973, A&A, 24, 337
 Shu, F., Najita, J., Ostriker, E., Wilken, F., Ruden, S., & Lizano, S. 1994, ApJ, 429, 781
 Voges, W., Atmanspacher, H., & Scheingraber, H. 1987, ApJ, 320, 794
 Wang, Y.-M. 1979, A&A, 74, 253